

Authentication of the Botanical and Geographical Origin of Honey by Front-Face Fluorescence Spectroscopy

KASPAR RUOFF,^{*,†,§} WERNER LUGINBÜHL,[†] RAPHAEL KÜNZLI,[§]
 STEFAN BOGDANOV,[†] JACQUES OLIVIER BOSSET,[†] KATHARINA VON DER OHE,[#]
 WERNER VON DER OHE,[#] AND RENATO AMADÒ[§]

Swiss Bee Research Centre/Agroscope Liebefeld-Posieux, Schwarzenburgstrasse 161, CH-3003 Berne, Switzerland; ETH Zurich, Institute of Food Science and Nutrition, ETH-Zentrum, Schmelzbergstrasse 9, CH-8092 Zurich, Switzerland; and Nds. Landesamt für Verbraucherschutz und Lebensmittelsicherheit, Institut für Bienenkunde Celle, Herzogin-Elionore-Allee 5, D-29221 Celle, Germany

Front-face fluorescence spectroscopy, directly applied on honey samples, was used for the authentication of 11 unifloral and polyfloral honey types ($n = 371$ samples) previously classified using traditional methods such as chemical, pollen, and sensory analysis. Excitation spectra (220–400 nm) were recorded with the emission measured at 420 nm. In addition, emission spectra were recorded between 290 and 500 nm (excitation at 270 nm) as well as between 330 and 550 nm (excitation at 310 nm). A total of four different spectral data sets were considered for data analysis. Chemometric evaluation of the spectra included principal component analysis and linear discriminant analysis; the error rates of the discriminant models were calculated by using Bayes' theorem. They ranged from <0.1% (polyfloral and chestnut honeys) to 9.9% (fir honeydew honey) by using single spectral data sets and from <0.1% (metcalfa honeydew, polyfloral, and chestnut honeys) to 7.5% (lime honey) by combining two data sets. This study indicates that front-face fluorescence spectroscopy is a promising technique for the authentication of the botanical origin of honey and may also be useful for the determination of the geographical origin within the same unifloral honey type.

KEYWORDS: Honey; unifloral; polyfloral; botanical origin; geographical origin; authenticity; adulteration; front-face fluorescence spectroscopy

INTRODUCTION

According to the Codex Alimentarius Standard (1) and the European Union Council Directive (2) relating to honey, the use of a botanical designation of honey is allowed if it originates predominately from the indicated floral source. Honey may also be designated by the name of a geographical region if it was produced within the area referred to (1, 2).

The vast majority of the honeys on the market contain significant nectar or honeydew contributions from several plant species and are therefore called polyfloral or multifloral honeys. Normally, they are just designated with the word "honey". Probably no honey produced by free-flying bees is purely unifloral. The term unifloral honey is used to describe honey in which the major part of the nectar or honeydew is derived from a single plant species. Honey composition, flavor, and color vary considerably depending on the botanical source it originates from (3).

The physical, chemical, and pollen analytical characteristics of the most important European unifloral honeys have been described in various papers (3–7). Contrary to the unifloral honeys, the polyfloral honeys do not exhibit distinct physical or chemical characteristic apart from a huge variability, which makes their authentication particularly difficult.

The interest in the production of unifloral honeys is caused by higher consumer preference for some honey types, generating a commercial concern of the beekeepers. The recent interest in the therapeutic or technological use of certain honey types may also contribute to the demand for a reliable determination of the botanical origin.

Botanical Origin. A number of new analytical techniques combined with multivariate data analysis have been proposed for the determination of the botanical origin of honey. They are, for example, based on physical and chemical measurands determined during quality control of honey (8, 9) or the former combined with the determination of mineral content (10), as well as carbohydrate composition (11), amino acid composition (12), mass spectrometry or metal oxide semiconductor based gas sensors (13, 14), differential scanning calorimetry (15),

* Address correspondence to this author at Inframiel, Herrenmatt 8, CH-5200 Brugg, Switzerland [telephone +41 (0)56 442 45 29; fax +41 (0)56 442 45 30; e-mail kaspar.ruoff@inframiel.ch].

[†] Swiss Bee Research Centre.

[§] ETH Zurich.

[#] Nds. Landesamt für Verbraucherschutz und Lebensmittelsicherheit.

pyrolysis mass spectrometry (16), and Raman (17) and near-infrared spectroscopy (18).

Many of the methods mentioned above allow one to clearly discriminate between several types of unifloral honeys, but none of these methods accounts for the polyfloral honeys that represent the majority of the honeys produced. This means that these methods may not be useful in analytical practice, as the great challenge in honey analytics is not to distinguish between several unifloral honey types but to discriminate the minority of unifloral honeys from the overwhelming majority of polyfloral honeys on the market. This also explains why until now none of these proposed methods are commonly used for the determination of the botanical origin of honey.

Only a single ion chromatographic method has been tested in the presence of polyfloral honeys and showed a potential to discriminate between several unifloral as well as polyfloral honey samples by first classifying the honey samples into two groups by color measurements (19). However, only very few samples were analyzed in this study, and it remains to be verified if this methodology is useful in analytical practice.

Currently, a reliable determination of the botanical and geographical origin can be achieved only by a global interpretation of sensory, pollen, and physicochemical analyses carried out by experts (4, 20, 21). However, the uncertainty related to the interpretation of pollen analytical results, originating from a number of different factors, demands the development of new analytical methods (22).

Geographical Origin. Pollen analysis is currently used to determine the geographical origin of honey as pollen in honey reflects the vegetation type where the nectar has been collected by the bees. In the past many analytical methods such as amino acid composition (23, 24), Raman spectroscopy (17), mineral content (25, 26), and sugar or mineral composition combined with common chemical quality control data (27–29) together with multivariate data evaluation have been proposed for the determination of the geographical origin.

Unfortunately, in most of the above quoted studies the botanical origin of the honey samples was not determined or the discrimination between the geographical origins was not verified on samples of the same botanical origin. Generally, the sample sets analyzed were small or limited to a small geographical area. The distinctions found are therefore rather due to differences of the vegetation type between the geographical regions and thus to the botanical origin of honey (30). A geographical discrimination will therefore be found when the differences are related to the vegetation type present in these areas.

As several analytical methods have to be used together for a reliable authentication of the botanical origin, such work is time-consuming and costly. Very specialized expertise is needed for the interpretation of the pollen spectrum used for the determination of the geographical origin of honey. Thus, there is a real need for new methods that allow a rapid and reproducible authentication of the botanical and geographical origin of honey at low cost (21, 31).

Fluorescence Spectroscopy. Compared to spectroscopic techniques based on absorption, fluorescence spectroscopy offers a 100–1000-fold higher sensitivity. It provides information on the presence of fluorescent molecules and their environment in inorganic and organic materials. In addition, front-face fluorescence spectroscopy allows an investigation of fluorophores in powders as well as in concentrated or opaque samples (32, 33).

Honey is known to contain fluorophores such as polyphenols (34–37) and amino acids (38, 39). Fluorescence spectroscopy should therefore be helpful for authenticating the botanical origin of honey. More detailed information on fluorescence spectroscopic applications to honey and other food can be found in our previous study, which already showed that front-face fluorescence spectroscopy is a promising approach for the determination of the botanical origin of honey (40).

The aim of the current work was to study the fluorescence spectroscopic characteristics of 11 honey types and to develop a rapid, low-cost, and reliable method for the authentication of unifloral and polyfloral honeys. As the physical and chemical characteristics of honey may be changed by adulteration, the potential of fluorescence spectroscopy was also studied on this subject. As minor nectar contributions from plant species other than the unifloral source may contribute to regional characteristics of unifloral honeys, the potential of fluorescence spectroscopy for the determination of the geographical origin of honey was studied as well.

MATERIALS AND METHODS

Sampling and Botanical Classification by Reference Methods.

A total of 371 honey samples produced between 1998 and 2004 were collected and stored at 4 °C until analysis. They originated predominantly from Switzerland (CH), but samples from Germany (D), Italy (I), Spain (E), France (F), Slovenia (SLO), and Denmark (DK) were also included.

To classify these honey samples corresponding to their botanical origin, the following measurands were determined according to the harmonized methods of the European Honey Commission (41): electrical conductivity, sugar composition, fructose/glucose ratio, pH value, free acidity, and proline content. Pollen analysis was carried out according to DIN 10760 (42, 43).

On the basis of these analytical results, the honey samples were assigned to one of the following 11 honey types, according to the criteria of Persano and Piro (3): acacia (*Robinia pseudoacacia*) (CH, $n = 14$; D, $n = 4$; F, $n = 3$); alpine rose (*Rhododendron* spp.) (CH, $n = 14$; I, $n = 5$); sweet chestnut (*Castanea sativa*) (CH, $n = 21$; I, $n = 5$; F, $n = 3$); rape (*Brassica napus* var. *oleifera*) (CH, $n = 22$); fir honeydew (*Abies* and *Picea* spp.) (CH, $n = 56$; D, $n = 63$; SLO, $n = 2$); oak honeydew (*Quercus* spp.) (E, $n = 8$); honeydew from *Metcalfa pruinosa* (I, $n = 14$); heather (*Calluna vulgaris*) (D, $n = 21$; DK, $n = 2$); lime (*Tilia* spp.) (CH, $n = 14$; D, $n = 9$; I, $n = 4$); dandelion (*Taraxacum* s.l.) (CH, $n = 10$; D, $n = 7$; I, $n = 2$); and polyfloral honeys (CH, $n = 68$). In the heterogeneous group of the polyfloral honeys, nectar or honeydew contributions from all of the above-mentioned sources were represented.

Adulterated Honeys. To evaluate the potential of fluorescence spectroscopy to detect beet sugar adulteration, an artificial honey was produced by feeding two colonies after the nectar flow, in autumn, with a sucrose solution of 62.5 g/100 g, generally used as winter feed for bee colonies in Switzerland. The sucrose solution was converted into artificial honey by the bees and left to ripen in the combs until extraction. To evaluate the possibility to detect honey adulteration by fluorescence spectroscopy, six chestnut and six acacia honey samples were adulterated with 50% of the artificial honey produced.

Fluorescence Spectroscopy. An aliquot of 20 g of the honey samples was liquefied at 55 °C for 8 h, allowed to cool to room temperature, and poured into a 1 cm quartz cuvette. The latter was placed into the sample holder of a Perkin-Elmer LS 50 B luminescence spectrometer (Perkin-Elmer, Beaconsfield, U.K.) equipped with a variable-angle front-surface accessory, with the incident angle of the excitation radiation set to 56°. Spectra were recorded at a scan rate of 150 nm/min and saved as ASCII textfiles. Instrumental artifacts were corrected in excitation using a rhodamine cell in the reference channel.

Method Development. To find additional wavelength ranges with specific emission or excitation for the honey types of interest in addition




Figure 1. Fluorescence spectra of different honey types.

to those already used in the preliminary study (40), the following ranges were studied. An excitation scan between 220 and 440 nm and recording of the fluorescence intensity at 420 and 490 nm was carried out. Six further emission scans were recorded with wavelengths between 220 and 600 nm, with excitation wavelengths being 210, 270, 310, 350, 390, and 440 nm, respectively. The following three instrumental settings yielded the most discriminating fluorescent spectra for the 10 types of unifloral honeys studied: excitation scan between 220 and 440 nm with the fluorescence emission measured at 420 nm (method A); using the excitation wavelengths of 270 and 310 nm, fluorescence emission spectra were recorded from 290 to 500 nm (method B) and from 330 to 550 nm (method C), respectively. The excitation slit width was set to 10 nm and the scan speed to 150 nm/min for all three methods. Two spectra were recorded using different aliquots of each sample. The spectra of the honey types studied are shown in **Figure 1**.

A control honey sample for the evaluation of instrumental stability and determination of the intermediate precision of the method was prepared by heating an acacia honey for 20 min up to 100 °C; the sample was then filtered to remove the pollen grains, partitioned into 2 mL glass vials, and then stored at -20 °C until analysis. The intermediate precision was determined by recording spectra of the control honey sample on 18 days of analysis within 1.5 months. The small coefficients of variation indicate that instrumental conditions were reasonably stable over the duration of the measurements (**Table 1**).

Processing of Spectra and Multivariate Analysis. The spectra were converted into the GRAMS spc-format (GRAMS/32 AI vs. 6.0, Thermo

Honey type

- Chestnut
- · - · Metcalfa honeydew
- · - · Dandelion
- Oak honeydew
- Fir honeydew
- - - - Heather
- · - · Acacia
- - - - Lime
- Alpine rose
- · - · Rape

Method

A: emission at 420 nm

B: excitation at 270 nm

C: excitation at 310 nm





Figure 2. Scatterplot of canonical discriminant scores from method A (for better legibility, the scores of only three honey types are displayed).

Galactic, Salem, NH) for more convenience in the visual examination and data reduction. It was found that a normalization of the spectra

Table 1. Repeatability and Intermediate Precision of the Three Fluorescence Spectroscopic Methods

| repeatability ($n = 6$) | method | | |
|-------------------------------------|--------|-------|-------|
| | A | B | C |
| av I_{\max} (au) ^a | 179.8 | 185.8 | 100.9 |
| reproducibility s_r (au) | 1.9 | 2.1 | 2.6 |
| coefficient of variation v_r (%) | 1.1 | 1.1 | 2.6 |
| repeatability limit (r) (au) | 5.4 | 5.9 | 7.5 |
| rel repeatability limit (%) | 3 | 3.2 | 7.4 |
| intermediate precision ($n = 18$) | | | |
| av I_{\max} (au) | 203 | 195 | 106 |
| lab reproducibility s_L (au) | 7.2 | 2.7 | 2.4 |
| rel lab reproducibility v_L (%) | 3.6 | 1.4 | 2.2 |

^a Arbitrary units.**Figure 3.** Scatterplot of canonical discriminant scores of lime honeys of different geographic origins (method A)


was not necessary and that the consideration of the fluorescence intensities can even improve the possibilities in discriminating the different unifloral honeys (40).

To avoid random noise resulting from instrumental effects, only the following spectral ranges were used for multivariate analysis: method A, 224–398 nm; method B, 290–500 nm; and method C, 333–547 nm. These ranges were also used for the combination of the spectra.

After elimination of spectral outliers, principal component analysis (PCA) was applied to eliminate the spectral collinearity and to reduce the number of variables to 20 PCs (using the PLSplus/IQ Add-on of GRAMS/32 AI vs. 5.09). This was performed separately for each type of spectra and each combination of different types of spectra.

In linear discriminant analysis (LDA), the 20 initial PCs were further reduced by backward elimination of PCs on the basis of their partial F values in the discriminant models (SYSTAT version 11, Systat Software Inc., Richmond, VA). The models were then optimized for maximum correct classification in jackknife classification. To account for the limited precision of single measurements, both spectra of each sample were used in the model of single types (A, B, and C) of spectra rather than the average. In the models using combined spectra, averaged spectra were used. The validation was carried out using spectra of one-third of the samples selected randomly and not present in the group of samples used to build the model.

Geographical Origin. The applicability of fluorescence spectroscopy for the determination of the geographical origin of honey was evaluated for the honey types when samples originating from different countries were available. The differences resulting from the geographic origin were studied within the groups of unifloral honeys by using MANOVA (SYSTAT version 11) as well as LDA and are visualized by plots of the canonical discriminant scores (Figures 3 and 4).

**Figure 4.** Scatterplot of the canonical discriminant score of fir honeydew honeys of German and Swiss provenance (method A).

RESULTS AND DISCUSSION

Repeatability. The repeatability of the three different methods was determined by a 6-fold measurement at the maximum intensity (I_{\max}) of an acacia honey. With coefficients of variation (v_r) between 1.1 and 2.6%, the methods showed a good repeatability, which stayed in the same range over 43 days while the intermediate precision was determined (Table 1).

Fluorescence Spectra of Different Honey Types. The recorded fluorescence spectra at three different excitation and emission wavelengths for the 10 unifloral honey types considered are displayed in Figure 1 (for better legibility, the spectra of alpine rose and rape honey are not shown in Figure 1A as they are visually very similar to that of fir honeydew honey). Every spectrum is typical for a given honey type. The spectra obtained by the different methods were recorded using different aliquots of the same sample.

Excitation spectra were scanned from 220 to 400 nm with the emission measured at 420 nm (method A, Figure 1A). For most honey types two maxima at ≈ 240 nm and between 340 and 360 nm, respectively, were observed, whereas lime honey exhibits its second maximum at ≈ 365 nm. Most of the honey types investigated had their intensity within the same order of magnitude except for chestnut honey, which showed a nearly 2-fold intensity at the maximum. Metcalfe honeydew honey is also characterized by a more intense fluorescence. Dandelion honey shows an additional shoulder at ≈ 300 nm.

For the spectra recorded using excitation at 270 nm (method B, Figure 1B), all honey types except chestnut, rape, and lime honeys exhibited broad and overlapping emission bands including at least two maxima located between 330 and 350 nm and between 400 and 440 nm, respectively. The very characteristic fluorescence spectrum of chestnut honey showed a much narrower band with two shoulders and a maximum at ≈ 380 nm. Rape and lime honeys showed both maxima at ≈ 350 nm, whereas the latter had a broader emission between 400 and 500 nm. Alpine rose honey showed a shoulder at ≈ 310 nm and a maximum at ≈ 340 nm. For heather, fir honeydew, dandelion, acacia, rape, and alpine rose honeys the intensities at the maxima ranged between 150 and 520 arbitrary units (au), whereas chestnut exhibited a considerably higher intensity of ≈ 800 au. The lowest intensity was detected for lime honey. However, the intensities were found to vary considerably within the honey types.

Using an excitation wavelength at 310 nm (method C, Figure 1C) the spectra of chestnut honey again clearly differed from

Table 2. Percentage of Correct Classification by Using Single Data Sets at Different Excitation and Emission Wavelengths and by Combining of the Data of the Methods (Jackknife Classification by the Leave One Out Method and Validation with Independent Samples)

| honey type | rates of correct classification by the different methods ^a (%) | | | | | | | |
|-------------------|---|------------|-----------|------------|-----------|------------|---|------------|
| | A | | B | | C | | combination of spectra from methods A and B | |
| | jackknife | validation | jackknife | validation | jackknife | validation | jackknife | validation |
| acacia | 95 | 100 | 90 | 79 | 85 | 75 | 90 | 100 |
| alpine rose | 87 | 100 | 50 | 80 | 63 | 50 | 93 | 100 |
| heather | 98 | 88 | 100 | 100 | 91 | 100 | 100 | 100 |
| chestnut | 97 | 100 | 96 | 100 | 96 | 100 | 96 | 100 |
| lime | 96 | 100 | 98 | 100 | 98 | 100 | 95 | 100 |
| dandelion | 100 | 100 | 97 | 100 | 100 | 100 | 100 | 100 |
| rape | 88 | 100 | 100 | 100 | 93 | 43 | 95 | 100 |
| fir honeydew | 92 | 86 | 91 | 84 | 84 | 76 | 96 | 97 |
| metcalfa honeydew | 93 | 80 | 100 | 100 | 100 | 75 | 92 | 100 |
| oak honeydew | 100 | | 100 | | 78 | | 100 | |
| polyfloral | 57 | 50 | 47 | 50 | 42 | 43 | 63 | 55 |
| av (weighted) | 87 | 85 | 84 | 83 | 80 | 73 | 90 | 91 |

^a Method A, excitation scanned 220 and 400 nm, emission measured at 420; method B, excitation at 270 nm, emission measured between 290 and 500 nm; method C, excitation at 310 nm, emission measured between 330 and 500 nm.

those of the other honey types investigated, especially by the 2-fold intensity compared to the others having a maximum at ≈ 380 nm. Most of the honey types exhibited a maximum at ≈ 400 nm and an intensity in the range from 100 to 200 au. Lime honey showed again the lowest intensity. Rape and acacia honeys were characterized by a shoulder at 365 nm. The maximum of the spectra of fir honeydew honeys was located at ≈ 355 and showed a shoulder at 420 nm. The band of the chestnut honey spectrum was narrower than that found by using method B and less intense but was, nevertheless, the most intense among the spectra recorded by using method C. The spectra of metcalfa honeydew honeys expressed a broad band with an intensity of ≈ 350 au, being thus the second most intense spectra.

It has been reported that chestnut honey, compared to the other honey types analyzed in this study, contains high amounts of hydroxycinnamates such as caffeic, *p*-coumaric, and ferulic acids as well as unidentified flavonoids (34, 44). Chestnut honey may also contain more phenylalanine than the other honey types analyzed in this study (39). The fluorescence of 2-aminoacetophenone, the main volatile component of chestnut honey, may also explain the characteristic spectra (45, 46).

Interestingly heather honey, commonly known to contain high amounts of phenolic compounds (47), does not show spectra of high fluorescence intensity compared to the other honey types. This may be due to scattering, reflection, and interference effects resulting from the numerous air bubbles present in heather honey.

LDA Applied to the Fluorescence Spectra. *Botanical Origin.* LDA was performed on the PCs of each type of spectra as well as on the combination of the two most significant types of spectra.

In the evaluation of single spectra the highest average classification rate (weighted according to the number of samples) of 85% in validation was obtained for method A (Table 2). The rates of correct classification were similar in both jackknife classification and validation, demonstrating that the models used were robust. Throughout the three methods studied, the classification rate for the polyfloral honeys was, at only 42–63%, very low. This can be explained by the lack of specific physical and chemical characteristics of this honey type. Thus, the polyfloral honeys are classified into the groups of unifloral honeys with the smallest Mahalanobis distance (Table 3).

For method A the lowest classification rate of 80% was observed for metcalfa honeydew honey. Twenty percent of the samples were misclassified as chestnut honey. This can be explained by the important nectar contribution of chestnut often present in metcalfa honeydew honeys. Due to the low number of samples ($n = 5$) used for validation, the 20% of misclassification arises from a single misclassified sample. In the validation step all samples of acacia, alpine rose, chestnut, lime, dandelion, and rape honeys were correctly classified. No validation was done for the oak honeydew honey due to the low number of samples available. In the jackknife classification some difficulties occurred in assigning alpine rose and acacia honeys (Table 3). Some samples of heather honey were also misclassified to rape and polyfloral honeys. Interestingly, a few samples of fir honeydew honey were classified as polyfloral or lime honeys. This could indicate that the value of 0.8 mS cm^{-1} in electrical conductivity is not always adequate to discriminate between polyfloral and honeydew honeys. Lime honeys very often contain some honeydew honey, which complicates their characterization.

Even though samples originated from different geographical origins, they were correctly classified according to their botanical origin. Irrespective of their geographical origin the fluorescent characteristics of honey from various botanical origins seem to be uniform, as samples from outside Switzerland group among the samples from Switzerland (Figure 2; for better legibility, the scores of only three different honey types are displayed).

The overall discriminating potential of method B is comparable to that of method A (Table 2). However, for the discrimination between alpine rose and acacia honeys more difficulties were encountered using method B than method A. In spite of the fact that the two groups were mingled, some samples of alpine rose honey were even misclassified as polyfloral honeys (data not shown).

The potential of method C for the classification of both unifloral and polyfloral honeys by using a single discriminant model was clearly inferior to that of methods A and B. Besides the difficulties already mentioned for alpine rose and acacia honeys, a considerable number of samples belonging to the groups of rape and honeydew honeys were not correctly classified in validation (Table 2).

To evaluate whether the rate of correct classification could be further increased by combining two of the most promising types of spectra, the ones of methods A and B were averaged

Table 3. Jackknife Classification and Validation Table for the Honey Samples Classified by LDA on the Spectra of Method A

| | jackknife classification rate for method A (%) | | | | | | | | | | |
|------------------------------------|--|----------------|-----------|-----------|-----------|------------|-----------|-----------------|----------------------|-----------------|------------|
| | acacia | alpine rose | heather | chestnut | lime | dandelion | rape | fir honeydew | metcalfa honeydew | oak honeydew | polyfloral |
| acacia (<i>n</i> = 21) | 95 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| alpine rose (<i>n</i> = 19) | 11 | 87 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| heather (<i>n</i> = 23) | 0 | 0 | 98 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| chestnut (<i>n</i> = 29) | 0 | 0 | 0 | 97 | 0 | 0 | 0 | 0 | 2 | 2 | 0 |
| lime (<i>n</i> = 26) | 0 | 0 | 0 | 0 | 96 | 0 | 0 | 0 | 0 | 0 | 4 |
| dandelion (<i>n</i> = 18) | 0 | 0 | 0 | 0 | 0 | 100 | 0 | 0 | 0 | 0 | 0 |
| rape (<i>n</i> = 24) | 0 | 0 | 0 | 0 | 0 | 0 | 88 | 0 | 0 | 0 | 12 |
| fir honeydew (<i>n</i> = 120) | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 92 | 0 | 0 | 8 |
| metcalfa honeydew (<i>n</i> = 14) | 0 | 0 | 0 | 7 | 0 | 0 | 0 | 0 | 93 | 0 | 0 |
| oak honeydew (<i>n</i> = 8) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 | 0 |
| polyfloral (<i>n</i> = 65) | 0 | 9 | 2 | 3 | 9 | 5 | 7 | 9 | 0 | 0 | 57 |

| | classification rate in validation for method A (%) | | | | | | | | | |
|-----------------------------------|--|----------------|-----------|------------|------------|------------|------------|-----------------|----------------------|------------|
| | acacia | alpine rose | heather | chestnut | lime | dandelion | rape | fir honeydew | metcalfa honeydew | polyfloral |
| acacia (<i>n</i> = 7) | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| alpine rose (<i>n</i> = 6) | 0 | 100 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| heather (<i>n</i> = 8) | 0 | 0 | 88 | 0 | 0 | 0 | 6 | 0 | 0 | 6 |
| chestnut (<i>n</i> = 10) | 0 | 0 | 0 | 100 | 0 | 0 | 0 | 0 | 0 | 0 |
| lime (<i>n</i> = 9) | 0 | 0 | 0 | 0 | 100 | 0 | 0 | 0 | 0 | 0 |
| dandelion (<i>n</i> = 6) | 0 | 0 | 0 | 0 | 0 | 100 | 0 | 0 | 0 | 0 |
| rape (<i>n</i> = 7) | 0 | 0 | 0 | 0 | 0 | 0 | 100 | 0 | 0 | 0 |
| fir honeydew (<i>n</i> = 40) | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 86 | 0 | 11 |
| metcalfa honeydew (<i>n</i> = 5) | 0 | 0 | 0 | 20 | 0 | 0 | 0 | 0 | 80 | 0 |
| polyfloral (<i>n</i> = 22) | 0 | 7 | 0 | 9 | 0 | 0 | 20 | 14 | 0 | 50 |

and concatenated for each sample. The rate of correct classification increased for alpine rose, fir, and spruce honeydew and even for polyfloral honeys compared to the results obtained by using the individual methods A and B (Table 2).

The classification tables revealed that polyfloral honeys were very often classified into the groups of the unifloral honeys, whereas the latter were rarely misclassified into the one of the polyfloral honeys. This observation led to the development of a two-step procedure. In the first step the sample was attributed to one of the 11 honey types considered using an overall discriminant model including all honey types. In the second step this classification was verified by using one or several two-group models consisting of a group formed by samples of a given unifloral honey versus a group called "non-unifloral", consisting of all the other samples. Each two-group model was separately built using LDA backward elimination and forward selection. For the verification of the classification by the first model, at least the two-group model of the corresponding honey type was used. In addition, one to six two-group models (entries in boldface type in Table 3) were used when a misclassification rate of >3% was calculated in jackknife classification or validation tables of the overall model.

The classification rates for the unifloral honeys in the two-group models were generally >90%, whereas the classification rate for the polyfloral honeys ranged between 48 and 75% (Table 4). However, as far as the polyfloral honeys are concerned, this is not very important, as we are principally interested in the authentication of unifloral honeys. The high rates of correct classification for both the unifloral and non-unifloral groups considered by the two-group models indicate that the botanical origin can be reliably determined according to this procedure. The respective error rates of this two-step procedure using methods A and B as well as the combination of the two former types of spectra were calculated by applying Bayes' theorem on the conditional probabilities of disjoint events.

Method A gave again the most promising results with an error probability (wrong classification of a sample of unknown botanical origin) of <5% for all honey types except for fir honeydew, for which it was 10% (Table 5). The error probabilities by using method B were higher for all honey types compared to those using method A except for the determination of metcalfa honeydew honey. By using the combination of the spectra of methods A and B, the error probability could be reduced to <5% (in validation) for the 11 honey types studied. It is interesting to note that the error probabilities of the honey types that express the highest variability in physical and chemical characteristics such as lime and fir honeydew honeys are the highest in fluorescence spectroscopy as well. This can be interpreted to indicate that fluorescence spectroscopy reproduces well the characteristics of classical criteria.

Geographical Origin. Differences in geographical origin were studied within the groups of samples of the same botanical origin when samples were available from at least two countries. Interestingly, a statistically significant difference was found by MANOVA between the geographical origins of all honey types studied (Table 6). The lime honey samples originating from Switzerland, Germany, and Italy formed groups in the plot of discriminant scores according to their geographical origin (Figure 3). The samples could also be correctly classified by LDA according to their geographical origin except for one Swiss sample that was classified to German provenance (data not shown). However, the classification according to geographical origin could be observed only within the groups of honeys of the same botanical origin. An LDA model of acacia, lime, dandelion, and fir honeydew honeys of German and Swiss origin failed to classify the samples according to their geographical provenance (Table 7). This clearly indicates that the characteristics resulting from the botanical source are considerably stronger than the geographical aspects. The sample set of the lime honeys was small; a larger sample set would possibly lead to a less pronounced difference. This may be illustrated on the

Table 4. Jackknife and Validation Tables for the Honey Samples Classified by the Two-Group Discriminant Models of Methods A and B and the Combination of These Spectra

| | jackknife classification | | | | validation | |
|---|--------------------------|----------------------------|---------------|----------------------------|------------|----------------------------|
| | unifloral | | non-unifloral | | unifloral | |
| | <i>n</i> | correct classification (%) | <i>n</i> | correct classification (%) | <i>n</i> | correct classification (%) |
| Method A | | | | | | |
| acacia | 21 | 100 | 343 | 96 | 7 | 100 |
| alpine rose | 19 | 100 | 345 | 90 | 6 | 83 |
| heather | 23 | 98 | 341 | 99 | 8 | 88 |
| chestnut | 29 | 97 | 335 | 99 | 10 | 90 |
| lime | 26 | 100 | 338 | 97 | 9 | 100 |
| dandelion | 18 | 100 | 346 | 98 | 6 | 100 |
| rape | 24 | 91 | 343 | 93 | 7 | 100 |
| fir honeydew | 120 | 95 | 244 | 93 | 40 | 88 |
| metcalfa honeydew | 14 | 100 | 350 | 100 | 5 | 100 |
| oak honeydew | 8 | 94 | 356 | 99 | | |
| polyfloral | 65 | 74 | 300 | 65 | 22 | 48 |
| Method B | | | | | | |
| acacia | 21 | 100 | 341 | 96 | 7 | 100 |
| alpine rose | 16 | 97 | 346 | 89 | 5 | 80 |
| heather | 23 | 100 | 340 | 95 | 8 | 100 |
| chestnut | 28 | 96 | 335 | 99 | 9 | 100 |
| lime | 26 | 100 | 336 | 95 | 8 | 100 |
| dandelion | 19 | 100 | 343 | 94 | 6 | 100 |
| rape | 22 | 100 | 340 | 98 | 7 | 100 |
| fir honeydew | 120 | 92 | 242 | 92 | 40 | 84 |
| metcalfa honeydew | 12 | 100 | 350 | 98 | 4 | 88 |
| oak honeydew | 8 | 100 | 354 | 95 | | |
| polyfloral | 68 | 65 | 294 | 72 | 22 | 75 |
| Combination of Spectra from Methods A and B | | | | | | |
| acacia | 20 | 100 | 327 | 97 | 7 | 100 |
| alpine rose | 15 | 100 | 332 | 91 | 7 | 100 |
| heather | 23 | 100 | 324 | 100 | 8 | 100 |
| chestnut | 27 | 96 | 320 | 100 | 9 | 100 |
| lime | 26 | 100 | 321 | 96 | 9 | 100 |
| dandelion | 19 | 95 | 328 | 98 | 6 | 100 |
| rape | 21 | 100 | 326 | 98 | 7 | 100 |
| fir honeydew | 117 | 97 | 230 | 93 | 39 | 100 |
| metcalfa honeydew | 12 | 100 | 335 | 98 | 4 | 100 |
| oak honeydew | 8 | 100 | 339 | 99 | | |
| polyfloral | 59 | 69 | 288 | 71 | 15 | 75 |

example of the fir honeydew honeys from Germany and Switzerland, for which a classification according to geographical origin was not possible (Figure 4). However, the samples of fir honeydew honeys originated from an area of ≈ 300 km in diameter belonging to Switzerland and Germany and therefore having very similar vegetation.

In future studies it should be verified if the geographical origin of honey could be determined by fluorescence spectroscopic

techniques on the basis of the minor contributions of accompanying flora that may be different in areas distant enough. The chemometric models should also be validated with samples of polyfloral provenance.

Adulteration by Feeding of Bees. The acacia and chestnut honey samples adulterated with as much as 50% of artificial honey did not show any comprehensible changes in the spectra compared to the pure samples in any of the three methods studied. Generally, the spectra of the adulterated samples remained in the range of the natural variation of the corresponding unifloral honeys. A detection of honey adulteration is therefore not possible except if the adulterant contains a characteristic fluorophore.

Conclusion. Although absolutely pure unifloral honeys do not exist, the definition of unifloral honey is in fact based on the points of view and the descriptions of different analysts. However, a consensus has been reached using the physical, chemical, and pollen analytical characteristics of the unifloral honeys considered as internationally recognized criteria already published (3–6).

Of capital importance is certainly to ensure a uniform honey quality that can be recognized by consumers preferring a given type of honey. Currently, the determination of the botanical origin of honey relies on the judgment of experienced experts who base their decision on the criteria of several analytical measurands. The challenge of new analytical methods that do not need such an expertise is to mathematically model and reproduce this decision-making process. As the definition of a unifloral honey is ultimately a matter of opinion, absolutely correct classification by chemometric models can therefore not be expected as these models are trained by uncertain sample sets as reference.

As the characteristic physical and chemical differences between unifloral and polyfloral honeys are small and only a very few compounds are specific to a given type of honey, the chemometric approach based on a fingerprint seems to be more promising than the search for individual marker compounds.

This study shows that front-face fluorescence spectroscopy combined with chemometrics offers a promising approach to the authentication of the botanical origin of honey and that the problems related to the determination of the polyfloral honeys can be overcome by the successive use of at least two mathematical models. The current results show that classical criteria commonly used for the determination of the botanical origin of honey can be very well reproduced by front-face fluorescence spectroscopy and chemometrics. Depending on the certainty needed, one may base the classification on the single spectra of type A or combine the spectra of methods A and B.

Table 5. Error Probabilities for the Classification of Unifloral and Polyfloral Honeys by the Different Methods

| honey type | error probability | | | | | |
|-------------------|-------------------|------------|-----------|------------|---|------------|
| | method A | | method B | | combination of spectra from methods A and B | |
| | jackknife | validation | jackknife | validation | jackknife | validation |
| acacia | 0.029 | 0.006 | 0.109 | 0.030 | 0.022 | 0.005 |
| alpine rose | 0.016 | 0.003 | 0.058 | 0.018 | 0.009 | 0.001 |
| heather | 0.044 | 0.013 | 0.050 | 0.051 | 0.003 | 0.003 |
| chestnut | 0.003 | $<10^{-3}$ | 0.034 | 0.053 | $<10^{-3}$ | $<10^{-3}$ |
| lime | 0.067 | 0.037 | 0.096 | 0.054 | 0.075 | 0.039 |
| dandelion | 0.037 | 0.008 | 0.075 | 0.072 | 0.021 | 0.019 |
| rape | 0.003 | 0.002 | 0.046 | 0.070 | 0.033 | 0.042 |
| fir honeydew | 0.088 | 0.099 | 0.107 | 0.090 | 0.047 | 0.045 |
| metcalfa honeydew | 0.040 | 0.004 | 0.004 | 0.002 | $<10^{-3}$ | $<10^{-3}$ |
| oak honeydew | 0.044 | | 0.050 | | 0.006 | |
| polyfloral | $<10^{-3}$ | $<10^{-3}$ | 0.034 | 0.031 | $<10^{-3}$ | $<10^{-3}$ |

Table 6. Results from MANOVA for the Geographical Origin of the Different Unifloral Honeys (Method A)

| honey type, country ^a | Wilks' λ | p |
|----------------------------------|------------------|-------------------|
| acacia (CH, D, F) | 0.009 | <10 ⁻³ |
| alpine rose (CH, I) | 0.027 | <10 ⁻³ |
| fir honeydew (CH, D) | 0.696 | <10 ⁻³ |
| chestnut (CH, F, I) | 0.001 | <10 ⁻³ |
| lime (CH, D, I) | 0.004 | <10 ⁻³ |
| dandelion (CH, D, I) | 0.023 | <10 ⁻³ |

^a Country codes: CH, Switzerland; D, Germany; F, France; I, Italy.

Table 7. Percentage of Correct Classification According to the Geographical Provenance by Using the Data Set of Method A

| | jackknife classification matrix ^a | | |
|-------------|--|---------|-------------|
| | Switzerland | Germany | correct (%) |
| Switzerland | 117 | 69 | 63 |
| Germany | 59 | 109 | 65 |
| total | 176 | 178 | 64 |

^a Jackknife classification by the leave one out method on samples from acacia, lime, dandelion, and honeydew honeys from spruce and fir.

Of course, the proposed fluorescence spectroscopic method needs a considerable amount of preliminary work to establish the chemometric models based on samples of known botanical origin. Once the classification models have been set, the technique enables a rapid determination of the botanical origin without particular sample preparation and special qualification of laboratory personnel. It remains to be tested by future studies if these models can be transferred from one instrument to another, as in infrared spectroscopy when normalized fluorescence spectra are used (40) or the instruments are calibrated with reference materials.

In addition, the present work clearly shows that fluorescence characteristics of honey are much more dependent on their botanical origin than on the geographical origin. Therefore, the former should be determined before a method is proposed for the determination of the geographical origin of honey. Such a method must be tested as well with samples of the same botanical origin.

ACKNOWLEDGMENT

The Istituto Nazionale di Apicoltura, Bologna, and Teresa Iglesias are acknowledged for the supply of honey samples from Italy and Spain, respectively.

LITERATURE CITED

- Codex Committee on Sugars, Revised Codex Standard for Honey. *Stand. Stand. Methods* **2001**, *11*, 1–7.
- Council Directive 2001/110/EC of 20 December 2001 relating to honey. *Off. J. Eur. Communities* **2002**, *L10*, 47–52.
- Persano Oddo, L.; Piro, R. Main European unifloral honeys: descriptive sheets. *Apidologie* **2004**, *35* (special issue), 38–81.
- Persano Oddo, L.; Piazza, M. G.; Sabatini, A. G.; Accorti, M. Characterization of unifloral honeys. *Apidologie* **1995**, *26*, 453–465.
- Piazza, M. G.; Persano Oddo, L. Bibliographical review of the main European unifloral honeys. *Apidologie* **2004**, *35* (special issue), 94–111.
- Moar, N. T. Pollen analysis of New Zealand honey. *N. Z. J. Agric. Res.* **1985**, *28*, 39–70.
- Crane, E.; Walker, P.; Day, R. *Directory of Important World Honey Sources. Nectar Plants*; International Bee Research Association: London, U.K., 1984.
- Devillers, J.; Morlot, M.; Pham-Delegue, M. H.; Dore, J. C. Classification of monofloral honeys based on their quality control data. *Food Chem.* **2004**, *86* (2), 305–312.
- Mateo, R.; Bosch-Reig, F. Classification of Spanish unifloral honeys by discriminant analysis of electrical conductivity, color, water content, sugars, and pH. *J. Agric. Food Chem.* **1998**, *46* (2), 393–400.
- Nalda, M. J. N.; Yague, J. L. B.; Calva, J. C. D.; Gomez, M. T. M. Classifying honeys from the Soria Province of Spain via multivariate analysis. *Anal. Bioanal. Chem.* **2005**, *382* (2), 311–319.
- Terrab, A.; Vega-Perez, J. M.; Diez, M. J.; Heredia, F. J. Characterisation of northwest Moroccan honeys by gas chromatographic–mass spectrometric analysis of their sugar components. *J. Sci. Food Agric.* **2002**, *82* (2), 179–185.
- Cotte, J. F.; Casabianca, H.; Giroud, B.; Albert, M.; Lheritier, J.; Grenier-Loustalot, M. F. Characterization of honey amino acid profiles using high-pressure liquid chromatography to control authenticity. *Anal. Bioanal. Chem.* **2004**, *378* (5), 1342–1350.
- Ampuero, S.; Bogdanov, S.; Bosset, J. O. Classification of unifloral honeys with an MS-based electronic nose using different sampling modes: SHS, SPME, and INDEX. *Eur. Food Res. Technol.* **2004**, *218*, 198–207.
- Benedetti, S.; Mannino, S.; Sabatini, A. G.; Marazzan, G. L. Electronic nose and neural network use for the classification of honey. *Apidologie* **2004**, *35*, 397–402.
- Cordella, C.; Faucon, J. P.; Cabrol-Bass, D.; Sbirrazzuoli, N. Application of DSC as a tool for honey floral species characterization and adulteration detection. *J. Therm. Anal. Calorim.* **2003**, *71*, 279–290.
- Radovic, B. S.; Goodacre, R.; Anklam, E. Contribution of pyrolysis–mass spectrometry (Py-MS) to authenticity testing of honey. *J. Anal. Pyrolysis* **2001**, *60* (1), 79–87.
- Goodacre, R.; Radovic, B. S.; Anklam, E. Progress toward the rapid nondestructive assessment of the floral origin of European honey using dispersive Raman spectroscopy. *Appl. Spectrosc.* **2002**, *56* (4), 521–527.
- Davies, A. M. C.; Radovic, B.; Fearn, T.; Anklam, E. A preliminary study on the characterisation of honey by near infrared spectroscopy. *J. Near Infrared Spectrosc.* **2002**, *10* (2), 121–135.
- Cordella, C. B.; Militao, J. S.; Clement, M. C.; Cabrol-Bass, D. Honey characterization and adulteration detection by pattern recognition applied on HPAEC-PAD profiles. I. Honey floral species characterization. *J. Agric. Food Chem.* **2003**, *51*, 3234–3242.
- Persano Oddo, L.; Bogdanov, S. Determination of botanical origin: problems and issues. *Apidologie* **2004**, *35* (special issue) 2–4.
- Bogdanov, S.; Ruoff, K.; Persano Oddo, L. Physico-chemical methods for the characterisation of unifloral honeys: a review. *Apidologie* **2004**, *35* (special issue), 4–17.
- Molan, P. C. The limitations of the methods of identifying the floral source of honeys. *Bee World* **1998**, *79* (2), 59–68.
- Gilbert, J.; Shepherd, M. J.; Wallwork, M. A.; Harris, R. G. Determination of the geographical origin of honeys by multivariate analysis of gas chromatographic data on their free amino acid content. *J. Apic. Res.* **1981**, *20* (2), 125–135.
- Davies, A. M. C. Amino acid analysis of honeys from eleven countries. *J. Apic. Res.* **1975**, *14* (1), 29–39.
- Latorre, M. J.; Pena, R.; Garcia, S.; Herrero, C. Authentication of Galician (NW Spain) honeys by multivariate techniques based on metal content data. *Analyst* **2000**, *125* (2), 307–312.
- Latorre, M. J.; Pena, R.; Pita, C.; Botana, A.; Garcia, S.; Herrero, C. Chemometric classification of honeys according to their type. II. Metal content data. *Food Chem.* **1999**, *66* (2), 263–268.

- (27) Gonzales Paramas, A. M.; Gomez Barez, J. A.; Garcia Villanova, R. J.; Rivas Pala, T.; Ardanuy Albajar, R.; Sanchez Sanchez, J. Geographical discrimination of honeys by using mineral composition and common chemical quality parameters. *J. Sci. Food Agric.* **2000**, *80*, 157–165.
- (28) Sanz, S.; Perez, C.; Herrera, A.; Sanz, M.; Juan, T. Application of a statistical approach to the classification of honey by geographic origin. *J. Sci. Food Agric.* **1995**, *69*, 135–140.
- (29) Gomez Barez, J. A.; Garcia Villanova, R. J.; Elvira Garcia, S.; Rivas Pala, T.; Gonzales Paramas, A. M.; Sanchez Sanchez, J. Geographical discrimination of honeys through the employment of sugar patterns and common chemical quality parameters. *Eur. Food Res. Technol.* **2000**, *210*, 437–444.
- (30) Cometto, P. M.; Faye, P. F.; Naranjo, R. D. D.; Rubio, M. A.; Aldao, M. A. J. Comparison of free amino acids profile in honey from three Argentinian regions. *J. Agric. Food Chem.* **2003**, *51*, 5079–5087.
- (31) Bogdanov, S.; Martin, P. Honey authenticity. *Mitt. Geb. Lebensmittelunters. Hyg.* **2002**, *93*, 232–254.
- (32) Parker, C. A. Apparatus and experimental methods. In *Photoluminescence of Solutions with Applications to Photochemistry and Analytical Chemistry*; Parker, C. A., Ed.; Elsevier: Amsterdam, The Netherlands, 1968; pp 128–302.
- (33) Genot, C.; Tonetti, F.; Montenay-Garestier, T.; Drapron, R. Front-face fluorescence applied to structural studies of proteins and lipid-protein interactions of visco-elastic food products. 1. Designing of front-face adaptor and validity of front-face fluorescence measurements. *Sci. Aliments.* **1992**, *12*, 199–212.
- (34) Amiot, M. J.; Aubert, S.; Gonnet, M.; Tacchini, M. Les composés phénoliques des miels: étude préliminaire sur l'identification et la quantification par familles. *Apidologie* **1989**, *20* (2), 115–125.
- (35) Andrade, P.; Ferreres, F.; Amaral, M. T. Analysis of honey phenolic acids by HPLC, its application to honey botanical characterization. *J. Liq. Chromat. Rel. Technol.* **1997**, *20* (14), 2281–2288.
- (36) Martos, I.; Ferreres, F.; Yao, L. H.; D'Arcy, B.; Caffin, N.; Tomás-Barberán, F. A. Flavonoids in monospecific *Eucalyptus* honeys from Australia. *J. Agric. Food Chem.* **2000**, *48*, 4744–4748.
- (37) Yao, L. H.; Jiang, Y. M.; D'Arcy, B.; Singanusong, R. T.; Datta, N.; Caffin, N.; Raymont, K. Quantitative high-performance liquid chromatography analyses of flavonoids in Australian *Eucalyptus* honeys. *J. Agric. Food Chem.* **2004**, *52*, 210–214.
- (38) Pirini, A.; Conte, L.; Francioso, O.; Lerkcer, G. Capillary gas chromatographic determination of free amino acids in honey as means of discrimination between botanical sources. *J. High Resolut. Chromatogr.* **1992**, *15*, 165–170.
- (39) Speer, K.; Montag, A. Verteilung freier Aminosäuren in Honigen. *Dtsch. Lebensm.-Rundsch.* **1986**, *82* (8), 248–253.
- (40) Ruoff, K.; Karoui, R.; Dufour, E.; Luginbühl, W.; Bosset, J. O.; Bogdanov, S.; Amadó, R. Authentication of the botanical origin of honey by front-face fluorescence spectroscopy. A preliminary study. *J. Agric. Food Chem.* **2005**, *53*, 1343–1347.
- (41) Bogdanov, S.; Martin, P.; Lüllmann, C. Harmonised methods of the European honey commission. *Apidologie* **1997**, (special issue), 1–59.
- (42) DIN. German Institute for Standardisation. *Analysis of Honey—Determination of the Relative Frequency of Pollen*; 2002.
- (43) von der Ohe, W.; Persano Oddo, L.; Piana, L.; Morlot, M.; Martin, P. Harmonised methods of melissopalynological analysis. *Apidologie* **2004**, *35* (special issue), 18–25.
- (44) Tomás-Barberán, F. A.; Martos, I.; Ferreres, F.; Radovic, B. S.; Anklam, E. HPLC flavonoid profiles as markers for the botanical origin of European unifloral honeys 34441. *J. Sci. Food Agric.* **2001**, *81*, 485–496.
- (45) Bonaga, G.; Giumanini, A. G. The volatile fraction of chestnut honey. *J. Apic. Res.* **1986**, *25* (2), 113–120.
- (46) Guyot, C.; Bouseta, A.; Scheirman, V.; Collin, S. Floral origin markers of chestnut and lime tree honeys. *J. Agric. Food Chem.* **1998**, *46*, 625–633.
- (47) Ferreres, F.; Andrade, P.; Tomás-Barberán, F. A. Flavonoids from Portuguese heather honey. *Z. Lebensm. Unters. Forsch.* **1994**, *1991*, 32–37.

Received for review March 12, 2006. Revised manuscript received June 20, 2006. Accepted June 22, 2006. We thank the Swiss Federal Office of Public Health, Narimpex AG, and the Swiss Beekeepers' Association for their financial support.

JF060697T